

Product Name: Overt Cockroach Gel Bait
 APVMA Approval No: 85164 / 112546



| | |
|-------------------------|---|
| Label Name: | Overt Cockroach Gel Bait |
| Signal Headings: | CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING |
| Constituent Statements: | ACTIVE CONSTITUENT: 6 g/kg INDOXACARB |
| Mode of Action: | GROUP 22A INSECTICIDE |
| Statement of Claims: | For the control of cockroaches in commercial, industrial and residential premises. |
| Net Contents: | Net Contents (5 g - 30 g) (single syringes) 10 g - 120 g (2 x 5 g to 4 x 30 g syringes) |
| Restrains: | RESTRAINTS DO NOT place gel on surfaces that are regularly washed. DO NOT apply this product in any areas easily accessible to children or animals. DO NOT allow product to directly contact humans, pets, exposed food, food preparation areas, food utensils, or food packing materials. |
| Directions for Use: | This section contains file attachment. |
| Other Limitations: | |
| Withholding Periods: | |

| | |
|---------------|--|
| Trade Advice: | |
|---------------|--|

| | |
|-----------------------|--|
| General Instructions: | <p>GENERAL INSTRUCTIONS</p> <p>To apply the gel, remove the cap on the nozzle and touch the tip to the surface to be treated and slowly depress plunger until sufficient gel is dispensed. Recap the dispenser after treatment is completed.</p> <p>Where maintenance treatments form part of the professional pest controllers contract it is recommended that cockroach activity be monitored where gel has been applied one month after initial treatment or at more regular intervals in areas of high infestation.</p> |
|-----------------------|--|

| | |
|---------------------|--|
| Resistance Warning: | <p>Insecticide Resistance Warning GROUP 22A INSECTICIDE</p> <p>Some naturally-occurring insect biotypes resistant to Overt Cockroach Gel Bait and other Group 22A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Overt Cockroach Gel Bait or other Group 22A insecticides are used repeatedly. The effectiveness of Overt Cockroach Gel Bait on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, PCT Holdings Pty Ltd accepts no liability for any losses that may result from the failure of Overt Cockroach Gel Bait to control resistant insects.</p> <p>Overt Cockroach Gel Bait may be subject to specific resistance management strategies. For further information contact PCT Holdings Pty Ltd or a local agricultural department agronomist.</p> |
|---------------------|--|

| | |
|--------------|--|
| Precautions: | |
|--------------|--|

| | |
|--------------|--|
| Protections: | <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</p> <p>DO NOT contaminate streams, rivers or waterways with the chemical or used containers.</p> |
|--------------|--|

| | |
|-----------------------|--|
| Storage and Disposal: | <p>STORAGE AND DISPOSAL</p> <p>Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.</p> |
|-----------------------|--|

| | |
|--------------------|--|
| Safety Directions: | <p>SAFETY DIRECTIONS</p> <p>Wash hands after use.</p> |
|--------------------|--|

| | |
|-------------------------|--|
| First Aid Instructions: | <p>FIRST AID</p> <p>If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone: Australia 13 11 26.</p> |
|-------------------------|--|

| | |
|---------------------|--|
| First Aid Warnings: | |
|---------------------|--|

| SITUATION | PEST | RATE |
|--|--|---|
| Domestic, commercial and public service buildings including: farm buildings, food processing establishments, factories, hospitals, hospitals, homes, kitchens, offices, restaurants, retail outlets, storerooms, vehicles and other situations which fall within these definitions. | German, American and other common cockroaches. | <p>Apply at least 2 spots per m². Increasing the number of spots (up to 5/m²) may be required for heavy infestations or if the American cockroach is the main pest.</p> <p>Recommended spot size is approximately 5 mm diameter (0.1 g) or the size of the spot indicated below:</p> <p style="text-align: center;">•</p> |
| CRITICAL COMMENTS | | |
| <p>Conduct a thorough inspection of the premises and then place gel in areas near cockroach harbourages or in sheltered areas where signs of cockroaches are evident.</p> <p>Replenish gel if previous application has been consumed or cockroaches remain a problem; increasing number of spots if necessary.</p> | | |

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION