POISON

KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Surefire Amine 625 Selective Herbicide

ACTIVE CONSTITUENT: 625 g/L 2,4-D present as the dimethylamine and diethanolamine salts



For the control of broadleaf weeds in fallow before direct drilling or sowing of cereals and pastures; and in cereal crops, pastures, sugarcane, peanuts, and non-agricultural areas as per the Directions for Use

This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops such as cotton, grapes, tomatoes, oilseed crops and ornamentals.

SPRAY DRIFT INSTRUCTIONS

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

PCT Holdings Pty Ltd

APVMA Approval No.: 82835/106882 (PCT Holdings Pty Ltd ABN 11 099 023 962) 5/74 Murdoch Circuit ACACIA RIDGE QLD 4110

http://pcti.com.au

CUSTOMER SERVICE FREECALL EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

DIRECTIONS FOR USE RESTRAINTS

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone tables below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply unless the wind speed is between 3 and 15 kilometres per hour at the application site during the time of application.

DO NOT apply if there are surface temperature inversion conditions present at the application site during the time of application. These conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise

Recognising a surface temperature inversion

A surface temperature inversion is likely to be present if:

- · Mist, fog, dew or a frost have occurred
- . Smoke or dust hangs in the air and moves sideways, just above the ground surface
- . Cumulus clouds that have built up during the day collapse towards evening
- Wind speed is constantly less than 11 km/hr in the evening and overnight
- · Cool off-slope breezes develop during the evening and overnight
- Distant sounds become clearer and easier to hear
- · Aromas become more distinct during the evening than during the day.

Spray timing

- Spray during the day wherever possible. Vertical mixing of the air makes surface temperature inversions unlikely and will reduce the risk of drift caused by surface temperature inversions.
- . There is a very low risk of surface temperature inversion when there is continuous overcast weather, with low and heavy cloud and/or wind speed remains above 11km/h for the whole period between sunset and sunrise.
- A lack of suitable weather conditions for spraying over extended periods is not an excuse for spraying in unsuitable conditions.

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions. DO NOT apply with spray droplets smaller than VERY COARSE spray droplets according to the ASAE S572.1 definition for standard nozzles.

DO NOT use if rain is likely within 6 hours.

Monitoring and record keeping

Users of this product MUST make an accurate written record of the details of each spray application within 24 hours following application and KEEP this record for a minimum of 2 years. The spray application details that must be recorded are: 1- date of use with start and finish times of application; 2- the specific location which must include address and paddock/s sprayed; 3- Product trade name (full name) of the product being used; 4- rate of application which must include the amount of product used per hectare and number of hectares applied to; 5- situation, crop or commodity to which the chemical was applied; 6- wind speed and direction during application; 7air temperature and relative humidity during application; 8- nozzle brand, model, size, type, and spray system pressure measured during application; 9- height of spay boom from ground; 10name and contact details of person applying this product (Additional record keeping and/or details may be required by the state or territory where this product is used).

Watch for changes in weather conditions. Stop spraying immediately if a surface temperature inversion occurs or if spraying conditions become unsuitable for any other reason.

ADVISORY FOR BOOM SPRAYER USE IN CEREALS, FALLOW AND PASTURE 1ST **OCTOBER TO 15 APRIL**

USE IN CEREALS, FALLOW AND PASTURES DURING THE PERIOD 3rd OCTOBER TO 15TH APRIL, IT IS ADVISED TO:-

USE NOZZLES THAT PRODUCE EXTREMELY COARSE (XC) TO ULTRA COARSE (UC)

USE HIGHER WATER RATES PER HA. TO GIVE BETTER EFFICACY.

USE SLOWER APPLICATION SPEEDS TO ALLOW OPERATORS TO LOWER BOOM

INCREASING DROPLET SIZE AND WATER RATES WHILE REDUCING APPLICATION SPEED WILL ASSIST IN MITIGATING OFF TARGET INVERSION DRIFT DURING SUMMER SPRAYING, EXTREMELY COARSE DROPLETS WILL PRODUCE < 3% DRIFTABLE DROPLETS.

BOOM SPRAYERS (ground application)

DO NOT apply by a boom sprayer unless the following requirements are met:

- . spray droplets not smaller than a VERY COARSE (VC) spray droplet size category (minimum XC between 3 October and 15 April - advisory)
- . boom heights 0.5 metres or lower above the target canopy (The higher of either the crop canopy or the targeted weeds)
- minimum distances between the application site and downwind sensitive aquatic and wetland areas including aquacultural ponds, surface streams and rivers (see Aquatic 'Downwind mandatory no-spray zone' section of the following table titled 'Buffer zones for boom sprayers') are observed.
- minimum distances between the application site and downwind sensitive crops, gardens, landscaping vegetation, protected native vegetation or protected animal habitat (see Terrestrial 'Downwind mandatory no-spray zone' section of the following table titled 'Buffer zones for boom sprayers') are observed. The buffer zones provide guidance but may not always be completely protective of all agricultural crops.

BUFFER ZONES FOR BOOM SPRAYERS:

Application rate (/ha)	Downwind manda	tory no spray zone
	Aquatic	Terrestrial
Dryland cropping: winter cereals	and fallows	AC.
Up to 1.2 L (750 g ae/ha)	10 metres	10 metres
Up to 1.4 L (880 g ae/ha)	15 metres	15 metres
Up to 1.7 L (1080 g ae/ha)	20 metres	20 metres
Dryland cropping: summer cerea	ls	
Up to 0.9 L (560 g ae/ha)	5 metres	0 metres
Tropical & subtropical uses: Sug	arcane	
Up to 1.8 L (1080 g ae/ha)	20 metres	20 metres
Up to 3.5 L (2220 g ae/ha)	35 metres	30 metres
Tropical & subtropical uses: Pea	nuts	
Up to 3.6 L (2250 g ae/ha)	35 metres	30 metres
Pasture		15.1
Up to 3.2 L (2000 g ae/ha)	30 metres	30 metres
Up to 4.4 L (2750 g ae/ha)	40 metres	35 metres
Up to 5.3 L (3300 g ae/ha)	45 metres	45 metres

Spray drift leaflet continued

Spray drift leaflet continued

Spray drift leaflet continued

AERIAL APPLICATION

DO NOT apply by aerial application unless the following requirements are met:

- spray droplets not smaller than a VERY COARSE (VC) spray droplet size category.
- · release heights 5 metres or lower above the target canopy
- minimum distances between the application site and downwind sensitive aquatic and wetland areas including aquacultural ponds, surface streams and rivers (see Aquatic 'Downwind mandatory no-spray zone' section of the following table titled 'Buffer zones for aircraft') are observed.
- minimum distances between the application site and downwind sensitive crops, gardens, landscaping vegetation, protected native vegetation or protected animal habitat (see Terrestrial 'Downwind mandatory no-spray zone' section of the following table titled 'Buffer zones for aircraft) are observed. The buffer zones provide guidance but may not always be completely protective of all agricultural crops.

BUFFER ZONES FOR AIRCRAFT: 3 metre release height or lower above the target canopy

Application rate (/ha)	D	Downwind mandatory no spray zone				
		Fixed wing		copter		
	Aquatic	Terrestrial	Aquatic	Terrestrial		
Dryland cropping: winter ce	reals and fallows		0			
Up to 1.2 L (750 g ae/ha)	75 metres	70 metres	70 metres	70 metres		
Up to 1.4 L (880 g ae/ha)	80 metres	80 metres	75 metres	75 metres		
Up to 1.7 L (1080 g ae/ha)	95 metres	90 metres	90 metres	85 metres		
Dryland cropping: summer	cereals					
Up to 0.9 L (560 g ae/ha)	60 metres	60 metres	60 metres	55 metres		
Tropical & subtropical uses	: Sugarcane					
Up to 3.5 L (2220 g ae/ha)	180 metres	170 metres	150 metres	140 metres		
Tropical & subtropical uses	: Peanuts	7				
Up to 3.5 L (2220 g ae/ha)	180 metres	170 metres	150 metres	140 metres		

BUFFER ZONES FOR AIRCRAFT: 5 metre release height or lower above the target canopy

Application rate (/ha)	Downwind mandatory no spray zone				
		d wing		copter	
	Aquatic	Terrestrial	Aquatic	Terrestrial	
Dryland cropping: winter ce	reals and fallows		10 -10		
Up to 1.2 L (750 g ae/ha)	130 metres	130 metres	120 metres	110 metres	
Up to 1.4 L (880 g ae/ha)	150 metres	150 metres	130 metres	120 metres	
Up to 1.7 L (1080 g ae/ha)	180 metres	170 metres	140 metres	140 metres	
Dryland cropping: summer	cereals	H WAS CONSTRUCT	The character		
Up to 0.9 L (560 g ae/ha)	110 metres	110 metres	100 metres	95 metres	
Tropical & subtropical uses	Sugarcane		10.		
Up to 3.5 L (2220 g ae/ha)	450 metres	400 metres	250 metres	225 metres	
Tropical & subtropical uses	Peanuts		-	-	
Up to 3.5 L (2220 g ae/ha)	450 metres	400 metres	250 metres	200 metres	

PASTURE APPLICATION BY AIR - 5.0 m RELEASE HEIGHT

Application rate 3330 g ae/ha, VERY COARSE droplet size, Aerial application

	Downwind no-spray zone		
Wind speed range at time of application	Fixed Wing	Helicopter	
From 3 to 7 kilometres per hour	600 metres	350 metres	
From 7 to 14 kilometres per hour	675 metres	375 metres	

Terrestrial protection

	Downwind no-spray zone		
Wind speed range at time of application	Fixed Wing	Helicopter	
From 3 to 7 kilometres per hour	575 metres	350 metres	
From 7 to 14 kilometres per hour	650 metres	350 metres	

Application rate 2750 g ae/ha, VERY COARSE droplet size, Aerial application Aquatic protection

	Downwind no-spray zone		
Wind speed range at time of application	Fixed Wing	Helicopter	
From 3 to 7 kilometres per hour	500 metres	300 metres	
From 7 to 14 kilometres per hour	550 metres	300 metres	

Terrestrial protection

	Downwind no-spray zone		
Wind speed range at time of application	Fixed Wing	Helicopter	
From 3 to 7 kilometres per hour	475 metres	275 metres	
From 7 to 14 kilometres per hour	525 metres	300 metres	

Application rate 2000 g ae/ha, VERY COARSE droplet size, Aerial application Aquatic protection

	Downwind no-spray zone		
Wind speed range at time of application	Fixed Wing	Helicopter	
From 3 to 7 kilometres per hour	375 metres	190 metres	
From 7 to 14 kilometres per hour	375 metres	220 metres	

Terrestrial protection

	Downwind no-spray zone		
Wind speed range at time of application	Fixed Wing	Helicopter	
From 3 to 7 kilometres per hour	350 metres	180 metres	
From 7 to 14 kilometres per hour	350 metres	210 metres	

PASTURE APPLICATION BY AIR- 3.0 m RELEASE HEIGHT

Application rate 3330 g ae/ha, VERY COARSE droplet size, Aerial application

BUILDING CONTROL CONTR	Downwind no-spray zone		
Wind speed range at time of application	Fixed Wing	Helicopter	
From 3 to 7 kilometres per hour	600 metres	350 metres	
From 7 to 14 kilometres per hour	675 metres	375 metres	

Terrestrial protection

Aquatic protection

	Downwind no-spray zone		
Wind speed range at time of application	Fixed Wing	Helicopter	
From 3 to 7 kilometres per hour	575 metres	350 metres	
From 7 to 14 kilometres per hour	650 metres	350 metres	

Application rate 2750 g ae/ha, VERY COARSE droplet size, Aerial application

	Downwind no-spray zone		
Wind speed range at time of application	Fixed Wing	Helicopter	
From 3 to 7 kilometres per hour	250 metres	150 metres	
From 7 to 14 kilometres per hour	250 metres	180 metres	

Terrestrial protection

Wind speed range at time of application	Downwind no-spray zone		
	Fixed Wing	Helicopter	
From 3 to 7 kilometres per hour	250 metres	140 metres	
From 7 to 14 kilometres per hour	250 metres	170 metres	

Application rate 2000 g ae/ha, VERY COARSE droplet size, Aerial application

Wind speed range at time of application	Downwind no-spray zone		
	Fixed Wing	Helicopter	
From 3 to 7 kilometres per hour	160 metres	90 metres	
From 7 to 14 kilometres per hour	160 metres	140 metres	

Terrestrial protection

Wind speed range at time of application	Downwind no-spray zone		
	Fixed Wing	Helicopter	
From 3 to 7 kilometres per hour	140 metres	85 metres	
From 7 to 14 kilometres per hour	150 metres	130 metres	